

# THE EFFECTS OF *Prosopis africana* PULP ON NUTRIENT DIGESTIBILITY, CARCASS COMPONENTS AND BLOOD COMPOSITION OF GROWING RABBIT

**\*Adamu, L.**

**Igwebuike, J. U.**

**Kwari, I. D.**

*Yobe State College of Agriculture,  
Gujba, Damaturu, Yobe State, Nigeria.*

**Aliyu, J.**

*Department of Animal Science,  
University of Maiduguri, Maiduguri, Borno State, Nigeria*

*\*E-mail: melelawan@yahoo.com*

## ABSTRACT

*A study was conducted to determine the effects of feeding *Prosopis africana* Pulp (PAP) on nutrient digestibility, haematological indices and carcass components of growing rabbits. The experiment which lasted 10 weeks consisted of thirty (30) rabbits of five to six weeks of age. They were randomly allotted to five dietary treatments in groups of six but caged individually. The PAP replaced maize weight for weight at levels of 0, 10, 20, 30 and 40% in diets 1 (control), 2, 3, 4 and 5 respectively. The formulated diets contained similar crude protein content of 18%. The apparent digestibility of nutrients, carcass parameters as well as haematological indices were not significantly different among all the treatments. These results indicate that *Prosopis africana* pulp can completely replace maize in the diets of growing rabbits without adverse effects on nutrient digestibility, carcass components and blood parameters.*

*Keywords: *Prosopis africana* pulp, rabbit feeding, digestibility, blood, carcass*

## INTRODUCTION

In many developing countries, one of the most important nutritional problems is inadequate intake of protein especially that of animal origin (Fielding, 1991). Animal protein contains more essential amino acids required to meet human nutritional needs than plant protein. FAO (1986) asserts that an average Nigerian consumed only 45.0g total protein and only 7.0g comes from animal protein, despite recommendation of 60 - 64g per individual daily out of which 27.3g should come from animal sources. The consequences are retarded growth, high incidence of kwashiorkor, high rate of child mortality, short life span and wide-spread protein - energy malnutrition (Adegbola, 1999). Production and multiplication of prolific monogastric animals and provision of adequate feeding appear to be some of the options for solving these problems. The ability of rabbits to thrive on forages and other similar feeds makes rabbit production comparatively cheaper (Aduku and Olukosi, 1990). Therefore, to

sustain interest, there is need to evaluate many feed resources, especially the legumes that are abundant during the dry season in the semi-arid areas of Nigeria.

*Prosopis africana* pods are among the earliest leguminous feeds known to man and still serve as a valuable source of carbohydrate and protein for many desert dwellers. Livestock also relish the pods which, in many species contain a sweet yellow pulp. It is used in ruminant feeding, but there is a dearth of information on its suitability in rabbit feeding. The objective of this study therefore was to assess the effect of *Prosopis africana* pulp on nutrient digestibility, carcass components and blood composition of growing rabbits.

## MATERIALS AND METHODS

The experimental site, experimental stock, management and experimental diets were similar to what was reported previously. At the end of week six of the experiment, three rabbits from each treatment were selected and placed individually in metabolism cages. Feed intake and faeces collected from each rabbit for five days were oven-dried and assayed for apparent digestibility of crude protein, ether extract, crude fibre, ash and nitrogen-free extract.

At week eight of the experiment, blood samples were also collected from three rabbits in each experiment for blood analyses. Blood samples were collected from the ear veins of the rabbits using sterile disposable syringe and needle. The rabbits were starved overnight and collection done the following morning at about 8.00a.m. in order to avoid excessive bleeding. Ethylenediamine tetracetic acid (EDTA) was used as an anticoagulant to prevent clotting of the blood collected for the haematological indices.

The Haematological indices which include Packed Cell Volume (PCV), Red Blood Cell (RBC) counts, White Blood Cell (WBC) counts and Haemoglobin (Hb) concentration were measured according to the methods expounded by Bush (1975). PCV, RBC, WBC and Hb were determined by micro haematocrit, improved Neubauer haemocytometer and cyanomethaemoglobin methods respectively. Mean Corpuscular Volume (MCV), Mean Corpuscular Haemoglobin (MCH) and Mean Corpuscular Haemoglobin Concentration (MCHC) were calculated according to standard formulae (Schalm, Jain and Carol, 1975).

Serum biochemical indices which include serum albumin, total protein, globulin, cholesterol, glucose, calcium (Ca) and Phosphorus (P) were measured according to methods described by Bush (1991). The serum glucose, urea and cholesterol were estimated by orthotoluidine, Diacetyl monoxime and colorimetric enzyme methods respectively. The total protein and serum albumin were determined by Biuret reactions (Bush, 1975).

At the end of week 10 of the study, three rabbits from each treatment were selected, weighed and then starved overnight (12 hours) but water was provided. The weights of the rabbit were recorded in the morning before slaughter. The dressed carcass, body components and organs were weighed and expressed as percentage of

slaughter weight. Dressed carcass and digestive tract length were also measured. The data collected were subjected to analysis of variance (ANOVA) using the completely randomized block design (Steel and Torrie, 1980). Means where significant ( $P < 0.05$ ) were separated using Least Significant Difference (LSD).

## RESULTS AND DISCUSSION

The mean apparent digestibility of dry matter, crude protein, crude fibre, ether extract and nitrogen free extract are presented on Table 1. There were no significant difference ( $P > 0.05$ ) in nutrient digestibility among all the treatment groups. This may be due to the similarity in the composition of the experimental diets, especially the protein contents. Although the digestibility of the nutrients are statistically similar, slightly higher digestibility was recorded in diet 1 (control) and this steadily decreased up to treatment 5. The progressive decrease in nutrient digestibility with increasing levels of PAP could not be attributed to high level of dietary fibre as observed by Butcher, Bryant, Machin, Owen, E. and Owen, J. (1981) and Onibi and Owa (1999) because the crude fibre in the diets (10.70 - 17.67) were within the normal range of 11.0 - 18.0% recommended in literature (Davidson and Spreadbury, 1975, Adegbola, Tibi and Asogwa, 1985; Igwebuiké, Abbas and Msheliza, 1998). It may however, be due to low feed intake and depressed nutrient intake and utilization arising from increasing tannin levels associated with the addition of PAP in the diets. Tannin is reported to depress nutrient digestibility and feed utilization (Mole, 1989).

On the other hand, Kassa, Abbas and Msheliza (1989) found a positive correlation between intake and digestibility of feeds in ruminants. Bohra and Ghosh (1980) and Ghol (1981) reported similar results with camels fed *Prosopis cineraria* pods. These observations are indications that the adverse consequences of tannins are more pronounced in non-ruminants than in ruminants (Jansman, 1993). Data on carcass characteristics are presented on table 2. There were no significant differences ( $P > 0.05$ ) in the slaughter weights among all the treatments. These results agree with those of Sankhyan, Wari and Narang (1991) and Igwebuiké, Alade and Anyi (1995) for growing rabbits. The dressing percentage values obtained in this study are lower than those reported by Garcia, Galvez and De-Blas (1993) who slaughtered rabbits of 2.0 - 2.5kg live weight and recorded dressing percentage of 58.0 and 60.0% respectively.

These may be attributed to the differences in slaughter weight since a positive correlation has been established between slaughter weight and dressing percentage (Garcia, Galvez and De-Blas, 1993). The components/organs shown on table 2 (head, skin, feet, heart, liver, lung, small intestine, large intestine, stomach, gullet, caecum, kidney and kidney fat) were weighed and expressed as a percentage of slaughter weight. They were not statistically different ( $P > 0.05$ ) among the treatment groups. The values recorded here compare favourably with those reported by Igwebuiké, Alade and Anyi (1995) who fed sorghum waste to growing rabbits in the same environment.

The length of gastro intestinal tract and dressed body weight showed no significant differences ( $P > 0.05$ ) among the different treatment groups. This indicates that the growth and development of these organs were not adversely affected by the various levels of PAP in the diets despite the variation in the protein, fibre and tannin levels of the diets. This is supported by the works of Aduku, Okoh, Njoku, Orjichie, Aganga and Dim (1986), Sankhyan, Wari and Narang (1991), Ijaiya and Fasanya (2004) who reported that the average dressing percentages and weights of body parts of rabbits fed different levels of protein and energy were not significantly ( $P > 0.05$ ) affected. The observation also agrees with the work of Zhao, Jorgensen and Eggum (1995) who reported that in rats, these components are hardly affected by the fibre levels of the diets. It is therefore concluded that the inclusion of PAP in diets of growing rabbits did not affect the proportionate development of the body components and organs.

### **Haematological and Erythrocytic indices**

The haematological and erythrocytic indices are presented in Table 3. The Packed Cell Volume (PCV) and Haemoglobin (Hb) concentration values were between 30% and 35% and 10.68 and 12.0g/100ml respectively. There were no significant differences ( $P > 0.05$ ) among the treatment means for PCV and Hb. All the values fall within the normal range of 31 to 50% and 8.0 to 15.97g/100ml for PCV and Hb respectively reported in literature (Schalm, Jain and Carol, 1975; CCAC, 1980). The normal and similar values obtained for all the treatment groups indicated nutritional adequacy of the various diets since abnormal values would have indicated mal - or undernutrition (Church, Judd, Young, Kebay and Kim, 1984). Hackbath, Buron and Schimansley (1983) found that there is a strong influence of diet on haematological traits with PCV and Hb being very strong indicators of nutritional status of animals. The PCV and Hb values obtained in this study are similar to the 29.50 to 36.5% and 10.40 to 12.6g/100ml reported by Alade, Kwaji and Igwebuiké (2001) for rabbits fed dry poultry waste. The values were however, slightly lower than the 38.0 to 41.25% and 10.30 to 11.05g/100ml PCV and Hb values respectively reported by Igwebuiké (2001) for growing rabbits.

The values for White Blood Cell (WBC) counts and red blood cell (RBC) counts are between 3.78 and 5.20 x 10<sup>3</sup>/ul, and 5.86 and 6.33 x 10<sup>6</sup>/ul respectively. There were no significant differences ( $P > 0.05$ ) between the treatment groups for either WBC or RBC counts. These values are within the normal range of 3.0 to 12.5 x 10<sup>3</sup>/ul and 3.0 to 7.73 x 10<sup>6</sup>/ul for WBC and RBC counts respectively (Schalm, Jain and Carol, 1975; CCAC, 1980). The slightly lower counts for WBC and RBC of rabbits fed diets 1, 4 and 5 compared to diets 2 and 3 may be attributed to the feed intake. Reduced feed intake normally leads to reduced nutrient intake which will include the iron needed for the formation of haemoglobin (Bush, 1975).

The WBC and RBC counts recorded in the study are lower than those reported by Igwebuiké (2001) who recorded the range of 5.28 to 9.05 x 10<sup>3</sup>/ul and 4.65 to

6.50 x 10<sup>6</sup>/ul for WBC and RBC counts respectively for growing rabbits fed *Acacia albida* pods. This may be attributed to the higher ambient temperature (30 - 42°C) during the period of the study. High ambient temperature (36 - 40°C) could lead to low WBC count in chickens (Kwari and Ubosi, 1991). Thermal stress has been shown to induce immuno-depression in birds (Subba-Rao and Glick, 1977) and this may apply to rabbits.

The values for Mean Corpuscular Volume (MCV) mean corpuscular Haemoglobin (MCH) and Mean Corpuscular Haemoglobin Concentration (MCHC) presented on table 3 were not significantly ( $P > 0.05$ ) different and were within the normal range of 60 - 73fl, 16 - 23pg and 26 - 34% for MVC, MCH and MCHC respectively reported by CCAC (1980). The normal values recorded in this study indicate that up to 40% PAP could be fed to growing rabbits without adverse effect on the blood components of growing rabbits.

The total protein, albumin and globulin are presented on table 4. There were no significant differences ( $P > 0.05$ ) among the treatment groups for these parameters. The values are within the normal ranges for 5.0 - 8.0g/dl, 3.13 - 3.80 g/dl and 2.27 to 2.97 g/dl for protein, albumin and globulin respectively as reported by CCAC (1980). Since these parameters are influenced by total protein intake (Birt and Schuldt, 1982), the values obtained in this study indicate nutritional adequacy of the dietary proteins. Abnormal serum albumin usually indicates an alteration of normal systematic protein utilization (Apata, 1990) and blood proteins are known to depend on the quality and quantity of dietary proteins (Awosanya, Joseph, Apata and Agboola, 1999). Therefore, the normal values obtained in this experiment indicate that the levels of PAP included in the diets had no adverse effect on serum protein level of the blood. The values for glucose and cholesterol ranged from 76.33 to 94.33mg/dl and 32.33 to 44.33mg/dl respectively and these were not significantly different among the treatment groups. They, however, fall within the normal reference values of 78 - 155mg/dl and 20.0 to 83.0 mg/dl respectively reported for normal healthy rabbits (CCAC, 1980; Onifade and Tewe, 1993). Since glucose and cholesterol levels were within the normal range, anorexia, diabetes, liver dysfunction and mal-absorption of fat which are symptoms of abnormal levels in the blood (Bush, 1991) are ruled out.

Serum calcium (Ca) and Phosphorus (P) obtained in this study were not significantly different ( $P > 0.05$ ) among the treatment groups. The values obtained compared favourably with the normal ranges of 5.6 - 12.7 mg/dl and 2.3 - 6.9 mg/dl for calcium and phosphorus levels of rabbits respectively (CCAC, 1980).

Since there was no sign of ill-health or anaemia observed in all the treatments, the diets might have met the minimum nutrient needs of the rabbits. Diets have very strong influence on haematological traits and serum biochemical indices (Hackbath et al., 1983). Abnormal values of these parameters would have indicated mal- or undernutrition (Duncan and Prasse, 1977 and Bush, 1975).

These results underline the usefulness of PAP for rabbit feeding. Haematological constituents reflect the physiological responsiveness of the animal to

its internal and external environments and these include feeds and feeding (Esonu et al., 2001). The normal blood indices obtained in this study shows that upto 40% PAP could be incorporated into the diets of growing rabbits without compromising the health status of the animals.

### CONCLUDING REMARK

The essence of protein intake is absolutely very significant for effective healthy living. This experiment evaluated the effect of feeding prosopis africana pulp on nutrient digestibility, carcass components and blood compositions of growing rabbit. The results underline the effectiveness of this diet for rabbit feeding. Since there was no sign of ill health or anemia observed in all the treatments, the diet was believed to have met the nutrients needs of the rabbits. The normal blood indices obtained in the study revealed that up to forty percent of the diet could be incorporated into the diet of growing rabbits without compromising their health status. Based on the results, it is therefore concluded that Prosopis africana pulp is sufficient feed substitute for growing rabbit and should be used by rabbit farmers to feed their rabbits.

**Table 1:** Mean apparent digestibility by rabbits fed graded levels of *Prosopis africana* pulp (PAP)

Nutrients	Diets/Treatments					SEM
	1	2	3	4	5	
Levels of PAP (%)	0	10	20	30	40	-
Dry matter	66.95	64.41	62.10	60.84	59.08	26.25NS
Crude protein	79.32	76.47	74.67	73.46	72.26	2.15NS
Crude fibre	38.41	37.68	37.57	35.98	34.49	19.99NS
Ether extract	70.27	68.08	66.19	65.92	61.11	10.00NS
Ash	60.09	59.08	57.59	56.95	54.87	16.69NS
Nitrogen-free Extract	89.89	86.59	93.36	79.65	78.92	21.33NS
SEM =	Standard Error of Means					
NS =	Not significant (P> 0.05)					

**Table 2:** Carcass data, weight and length of selected organs of rabbits fed graded levels of *Prosopis africana* pulp (PAP)

Parameters	Diets/Treatment groups					SEM
	1	2	3	4	5	
Level of PAP (%)	0	10	20	30	40	-
No. of rabbits						
Slaughtered	3	3	3	3	3	-
Slaughter weight (g)	1483.00	1300.00	1307.00	1250.00	1033.00	1100.0NS
Wt. of dressed carcass (g)	689.9	682.71	675.33	637.36	670.54	55.28NS
Dressing percentage (%)	46.96	47.57	46.94	45.64	44.90	10.20NS
Body Components as Percentage of Slaughter weight (%)						
Head	7.668	8.329	7.856	8.513	9.624	0.57NS
Skin	4.818	6.800	6.186	6.254	6.429	1.095NS
Tail	0.335	0.305	0.297	0.243	0.325	0.009NS
Feet	2.122	2.274	2.329	2.585	2.603	0.177NS
Heart	0.211	0.229	0.218	0.260	0.178	0.001NS
Liver	2.796	2.826	2.581	2.458	2.827	0.171NS

Lung	0.449	0.465	0.458	0.576	0.641	0.022NS
Spleen	0.025	0.024	0.032	0.028	0.031	0.002NS
Kidneys	0.468	0.500	0.495	0.491	0.605	0.006NS+
Oesophagus	0.823	0.824	0.856	0.831	0.793	0.005NS
Stomach	5.628	5.106	6.538	6.569	7.902	1.513NS
Small intestine	3.660	3.688	3.778	3.599	3.568	0.035NS
Large intestine	5.100	6.075	6.172	5.061	4.997	0.89NS
Caecum	0.831	0.841	8.873	0.712	0.790	0.004NS
Kidney fat	0.507	0.623	0.681	0.281	0.39	0.062NS
Length of Gastro-intestinal tract (CM)						
Oesophagus	10.30	10.33	10.50	10.167	10.00	0.718NS
Stomach	9.00	9.167	9.267	8.33	7.67	0.48NS
Small intestine	233.50	234.33	235.68	233.33	230.33	209.00NS
Large intestine	143.20	143.33	144.33	143.00	142.00	5.945NS
Caecum	9.667	9.667	9.00	11.50	9.00	2.77NS
Body length	31.00	29.23	29.17	29.33	28.17	1.58NS
PAP	=	<i>Prosopis africana</i> Pulp				
SEM	=	Standard error of means				
NS	=	Not significant ( $P > 0.05$ )				

**Table 3:** Haematological indices of rabbits fed graded levels of *Prosopis africana* pulp (PAP)

Indices	Diets/Treatments					SEM
	1	2	3	4	5	
Levels of PAP (%)	0	10	20	30	40	-
Haematocrit (PCV %)	34.35	35.00	35.00	33.00	33.00	6.73NS
Haemoglobin (Hb g/100ml)	11.33	12.0	12.0	11.68	10.68	1.23NS
WBC (x 10 <sup>3</sup> /mm <sup>3</sup> )	4.28	4.70	5.20	3.88	3.78	1.28NS
RBC (x 10 <sup>6</sup> /mm <sup>3</sup> )	5.73	6.31	6.31	6.01	5.86	0.71NS
MCV (fl)	54.13	55.49	57.85	55.62	50.94	7.83NS
MCH (pg)	19.69	19.08	19.88	19.68	18.17	8.47NS
MCHC (%)	43.95	34.29	34.28	36.11	35.45	0.79NS
NS	=	Not significant ( $P > 0.05$ )				
PCV	=	Packed Cell Volume				
WBC	=	White Blood Cell				
RBC	=	Red Blood Cell				
MCV	=	Mean Corpuscular Volume				
MCH	=	Mean Corpuscular Haemoglobin				
MCHC	=	Mean Corpuscular Haemoglobin Concentration				
SEM	=	Standard Error of Means				

**Table 4:** Serum biochemical indices of rabbits fed graded levels of *Prosopis africana* pulp (PAP)

Indices	Diets/Treatments					SEM
	1	2	3	4	5	
Levels of PAP (%)	0	10	20	30	40	-
Total protein (g/dl)	6.58	6.30	6.73	6.10	5.70	1.19NS
Albumin (g/dl)	3.68	3.33	3.80	3.23	3.43	0.08NS
Globulin (g/dl)	2.90	2.97	2.93	2.97	2.22	0.15NS
Glucose (mg/dl)	86.67	85.33	94.33	86.67	76.33	13.98NS
Cholesterol (mg/dl)	40.33	39.0	44.33	32.33	40.0	7.15NS
Calcium (mg/dl)	6.33	6.33	6.42	6.33	6.31	0.003NS
Phosphorus (mg/dl)	2.47	2.40	3.10	2.37	2.13	0.11NS
SEM	=	Standard error of means				
NS	=	Not significant ( $P > 0.05$ )				

## REFERENCES

- Adegbola, T. A., Tibi, E. U. and Asogwa, D. C.** (1985). Feed intake and digestibility of rabbit on all-forage, forage plus concentrate and all concentrate diets. *Journal of Animal Production Research*, 5(2), 185 - 191.
- Adegbola, T. A.** (1999). Livestock production toward alleviation of poverty and malnutrition in families. In Proceedings of International Seminar on Promoting sustainable small scale livestock and poverty in Rural land Sub-urban families in Nigeria. March, 7 - 11, 1999. Abubakar Tafawa Balewa University, Bauchi, Nigeria. 1981 - 193.
- Aduku A. O., Okoh P. N., Njoku P. C., Orjichie E.A., Aganga A. A. and Dim N. I.** (1986). Evaluation of cowpea (*Vigna unguiculata*) and peanut (*Arachis hypogea*) haulms as feed stuffs for weanling rabbits in a tropical environment (Nigeria). *Journal of Applied Rabbit Research*, 91, 178 - 179.
- Aduku, O. A. and Olukosi, J. D.** (1990). Rabbit Management in the tropics: Production, Processing, Utilization, Marketing, Economics, Practical Training, Research and Future Prospects. Abuja: Living Book Series, G. U. Publishers, FCT. 1 - 105.
- Alade, N. K., Kwaji, D. T. and Igwebuik, J. U.** (2001). Growth performance and blood constituents of rabbits fed graded levels of poultry waste. *Annals of Borno*, L17/18: 217 - 225.
- Apata, D. F.** (1990). Biochemical, Nutritional and Toxicological assessment of some tropical legume seeds. Ph. D. Thesis, University of Ibadan, Ibadan, Nigeria.
- Awosanya, B; Joseph, J. K., Apata, D.F. and Agboola, M. A.** (1999). Performance, blood chemistry and carcass quality attributes of rabbits fed raw and processed *peuraria* seed meal. *Tropical Journal of Animal Science*, 2 (2), 89 - 96.
- Birth, D. F. and Schuldt, G. H.** (1982). Effects of the different source and level of protein fed to Syrian hamsters on growth, protein utilization and selected blood proteins. *Laboratory Animal Science*, 32, 617 - 630.
- Bohra, M. C. and Ghosh, P. K.** (1980). *The nutritive value and digestibility of Prosopis species in the Indian desert*. New Delhi: CAZRI - ICAR, India.
- Bush, B. M.** (1975). *Veterinary Laboratory Manual*. London: William Heinemann Medical Book Ltd., U.K.
- Bush, B. M.** (1991). Interpretation of Laboratory Results for Small Animal Clinicians. United Kingdom: Blackwell Scientific Publications.
- Butcher, C., Bryant, M. J., Machin, D. M., Owen, E. and Owen, J. E.** (1981). The effect of metabolic energy concentration on the performance in digestibility of growing rabbits. *Tropical Animal Production*, 6, 93 - 100.
- Canadian Council on Animal Care (CCAC)** (1980). Guide to the Care and Use of Experimental Animals. Vol. 1. Canadian Council on Animal Care, Ottawa, Ontario, Canada. 85 - 90.
- Church J. P., Judd J. T., Young C. W., Kebay T. L. and Kim W. W.** (1984). Relationship among dietary constituents and specific serum clinical components of subjects eating self-selected diets. *American Journal of Clinical Nutrient*, 40, 1338 - 1344.
- Davidson, W. J. and Spreadbury, D.** (1975). Nutrition of New Zealand White Rabbit. Proceedings of Nutrition Society, 34, 75 - 83.
- Duncan, R. J. and Prasse, K. W.** (1977). *Veterinary Laboratory Medicine. Clinical Pathology*. Iowa: The Iowa State University Press, Ames., U. S. A. 243.
- Esonu B. O., Emenalom O. O., Udedidbie A. B. I., Herbert U., Ekpor C. F., Okoli I. C. and Iheukwumere F. C.** (2001). Performance and blood chemistry of weaner pigs fed raw mucuna bean (Velvet bean) meal. *Tropical Animal Production Investigation*, 4, 49 - 54.
- FAO** (1986). The state of Food and Agriculture. Paper presented in Food and Agriculture Organisation (FAO), Summit, Geneva, 1986.



- Fielding, D.** (1991). Rabbit. London and Basingstoke: The Macmillan Press Ltd.
- Garcia G., Galvez J. F. and De-Blas J. C.** (1993). Effects of substitution of sugar beet pulp for barley in diets for finishing rabbit on growth performance and on energy and nitrogen efficiency. *Journal of Animal Science*, 71, 1823 - 1830.
- Ghol, B.** (1981). Tropical Feeds: Feed information summaries and Nutritional value. FAO Animal production and Health series. No. 12. Food and Agriculture Organisation, Rome, Italy. 121 - 529.
- Hackbarth, H; Buron, K. and Schimansley, C.** (1983). Strain differences in inbred rats: Influence of strain and diet on haematological traits. *Laboratory Animal*, 17, 7 - 19.
- Igwebuike J. U., Alade N. K. and Anyi H. D.** (1995). Effect of feeding graded levels of sorghum waste on the performance and organ weights of growing rabbits. *East Africa Journal of Agricultural Forestry*, 60 (40), 193 - 200.
- Igwebuike J. U., Abbas I. and Msheliza N. K. A.** (1998). Effect of feeding graded levels of sorghum waste on the nutrient and mineral utilization of growing rabbits. *Reserch Journal of Science*, 4(1 & 2), 46 - 56.
- Igwebuike, J. U.** (2001). Utilization of Acacia pod, (*Acacia albida* Del.) for Rabbit feeding. Ph. D. Thesis submitted to University of Agriculture, Makurdi, Nigeria.
- Ijaiya, T. A. and Fasanya, O. O. A.** (2004). Effects of varying levels of dietary protein on the carcass characteristics of growing rabbits. *Nigerian Journal of Animal Production*, 31 (2), 207 - 210.
- Jansman, A. J. M.** (1993). Tannins in feedstuff for simple-stomached animals. *Nutrition Research Review*, 6, 209 - 236.
- Kassa W., Thwaites C. J., Jianke X. and Farrel D. J.** (1989). Rice bran in the diets of rabbits growth at 22oC and 30oC. *Journal of Applied Rabbit Research*, 12, 75 - 77.
- Kwari, I. D. and Ubosi, C. O.** (1991). Comparison of three housing systems for growing broiler chickens in hot dry climate 2. Effect of ambient temperature on blood components. *Journal of Agricultural Science Technology*, 1 (2), 123 - 125.
- Mole, S.** (1989). Polyphenolics and the nutritional ecology of herbivores. In P. R. Cheeke ed) Toxicants of plant origin. IV. Phenolics Boca Raton Florida: CRC Press USA 191 - 223.
- Onibi, G. E. and Owa, B. O.** (1999). Influence of period of provision of commercial pellets and forage (*Aspilla africana*) on the growth performance and economics of production of rabbits. In Kolade J., Awosanya, B, Apata, D.d F. and Balewu, M. A. (eds.) Enhancing Livestock Production in Nigeria. Proceedings of the 26th Annual Conference of Nigerian Society of Agricultural Practitioners, Kwara Hotels, Ilorin, 21 - 25th March, 1999. Pp. 151 - 152.
- Onifade, A. A. and Tewe, O. O.** (1993). Alternative tropical energy feed in rabbit diets; growth performance, diets digestibility and blood composition. *World Rabbit Science*, 1(1), 17 - 24.
- Sankhyan, S. T.; Wari, S. P. and Narang, M. P.** (1991). Effect of dietary protein and energy levels on the carcass characteristics of rabbits. *Journal of Applied Rabbit Research*, 14, 54 - 56.
- Schalm, O. W., Jain, N. C. and Carol, E.** (1975). *Veterinary Haematology* (3rd edn.) Philadelphia: Lea and Febiger, U. S. A.
- Steel, R. G. D. and Torrie, J. H.** (1980). *Principles and Procedures of Statistics. A Biometrical Approach* (2nd edn). New York McGraw Hill Book Co., USA.
- Subba - Rao, D. S. V. and B. Glick** (1977). Immuno-suppressive action of heat in chickens. *Proceedings of the Society for Experimental biology and Medicine*, 131, 445 - 465.
- Zhao X., Jorgensen H. and Eggum B. D.** (1995). The influence of dietary fibre on body composition, visceral organ weight, digestibility and energy in rats housed in different thermal environment. *British Journal of Nutrition*, 73, 687 -699.